

Improving the thermostability and catalytic efficiency of a GH10 xylanase by rational design for optimizing the recycle of agricultural waste

Yifan Zhao¹, Chao Zhu¹, Yufeng Zhao¹, Aoxue Li¹, Leyun Yang^{1,2}, Chunying Feng^{1,2}, Shuai You^{1,2*}, Jun Wang^{1,2*}

¹ Jiangsu Key Laboratory of Sericultural Biology and Biotechnology, School of Biotechnology, Jiangsu University of Science and Technology, Zhenjiang 212100, China;

² Key Laboratory of Silkworm and Mulberry Genetic Improvement, Ministry of Agriculture and Rural Affairs, The Sericultural Research Institute, Chinese Academy of Agricultural Sciences, Zhenjiang 212100, China.

Keywords: GH10 Xylanase, Degradation of agricultural lignocellulosic wastes, Protein engineering

Presenting author email: youshuai@just.edu.cn.

Endo- β -1,4-xylanases (EC 3.2.1.8) catalyze the hydrolysis of β -1,4-glycosidic bonds in heteroxylans, which are major components of hemicellulose in the biosphere (Prade R. 1995). The enzyme has considerable potential in many applications, such as deinking of waste paper, improving the digestibility of animal feed, bread making, converting of plant biomass into biofuels, and producing prebiotics (Tony et al., 2010).

Xylanases are mainly distributed in glycoside hydrolase (GH) families 5, 7, 8, 10, 11, 30, and 43. Current research focuses on GH10 and GH11 family xylanase. The GH10 xylanases has better thermal stability (Beaugrand et al., 2004), but lower catalytic efficiency, which limits its industrial application. Therefore, obtaining xylanase with high thermocatalytic activity is very important for industrial production.

At present, protein engineering is widely used in the improvement of enzyme molecules, that is, by modifying or modifying genes or the protein itself to change the structure of the protein to achieve the transformation of enzyme functions. Protein engineering is mainly used for the design and modification of enzyme properties such as thermal stability, catalytic efficiency, substrate specificity and extreme environmental tolerance. The main methods involved are directed evolution, rational design and semi-rational design. Among them, rational design is a quick and effective means of transformation, and its commonly used methods mainly include module replacement and site-specific mutation. This method significantly improves the thermostability of the xylanase XynA derived from *Thermoascus aurantiacus* (Abou-Hachem et al., 2002). Wang et al. optimized the catalytic channel of TlXyn10A through rational design, which increased the specific activity by 40% (Wang et al., 2016).

Based on the relationship between structure and function, in this study, the GH10 xylanase (GtXyn10) from *Gloeophyllum trabeum* via site-directed mutagenesis to improve the catalytic performance. Through catalytic amino acid sites and substrate binding sites were predicted by hotspot wizard, three locations, G29L, H32K, and Q99A were identified as the contributors to the whole enzyme molecule. The mutants were produced and compared with the wild-type (WT). Using beechwood xylan as substrate, the specific activities of mutants G29L, H32K and Q99A were 1520 U/mg, 1830 U/mg and 1720 U/mg, respectively, which were higher than that of wild-type GtXyn10 (1060U/mg) were increased by 43%, 73% and 62%, the catalytic efficiency was 300 mL/s·mg, 610 mL/s·mg and 360 mL/s·mg respectively, compared with wild-type GtXyn10 (210 mL/s·mg), increased by 43%, 1.9 times and 71% respectively (Table 1).

The optimum temperature and optimum pH have no significant changes compared to the wild type. In terms of thermal stability (Figure 1), mutants G29L, H32K and Q99A were treated at 75 °C for 30 min, and the remaining enzyme activities were 45%, 64% and 28%, respectively. Compared with wild-type enzyme GtXyn10 (62%), the thermal stability of mutants G29L and H32K did not change significantly, while the thermal stability of mutant Q99A decreased significantly.

Table 1. Kinetic parameters and specific activity of GtXyn10 and its mutants towards beechwood xylan.

Enzymes	K_m (mg/mL)	V_{max} ($\mu\text{mol}/\text{min}\cdot\text{mg}$)	k_{cat}/K_m (mL/s \cdot mg)	Specific activity (U/mg)
GtXyn10	4.2 ± 0.41	1450 ± 179	210 ± 25	1060 ± 95
G29L	4.3 ± 0.38	2080 ± 301	300 ± 41	1520 ± 132
H32K	2.4 ± 0.15	2380 ± 229	610 ± 77	1830 ± 199
Q99A	3.8 ± 0.27	2250 ± 218	360 ± 49	1720 ± 183




Fig. 1 Relative activity analysis of recombinant xylanase mutants and wild type

A, the optimal pH; B, the pH stability; C, the optimal temperature; D, thermal stability at 75°C.

Acknowledgements

This work was supported by the the Jiangsu Agricultural Science and Technology Innovation Fund (CX(22)3034), the National Natural Science Foundation of China (21978121), China Agriculture Research System of MOF and MARA, and Key Projects of International Scientific and Technological Innovation Cooperation of National Key R&D Program of China (2021YFE0111100).

References:

- [1] Abou-Hachem, M., Karlsson, E.N., Simpson, P.J., Linse, S., Sellers, P., Williamson, M.P., Jamieson, S.J., Gilbert, H.J., Bolam, D.N., Holst, O., 2002. Calcium binding and thermostability of carbohydrate binding module CBM4-2 of Xyn10A from *Rhodothermus marinus*. *Biochemistry-US*, 41(18): 5720-5729.
- [2] Beaugrand, J., Chambat, G., Wong, V., Goubet, F., Rémond, C., Pa S, G., Benamrouche, S., Deb Eire, P., Donohue, M.O., Chabbert, B., 2004. Impact and efficiency of GH10 and GH11 thermostable endoxylanases on wheat bran and alkali-extractable arabinoxylans. *Carbohydr. Res.*, 339(15): 2529-2540.
- [3] Tony, C., Charles, G., Georges, F., 2010. Xylanases, xylanase families and extremophilic xylanases. *FEMS Microbiol. Rev.*, 29(1): 3-23.
- [4] Wang, X., Huang, H., Xie, X., Ma, R., Bai, Y., Zheng, F., You, S., Zhang, B., Xie, H., Yao, B., 2016. Improvement of the catalytic performance of a hyperthermostable GH10 xylanase from *Talaromyces leycettanus* JCM12802. *Bioresour. Technol.*, 222(1): 277-284.
- [5] Prade R., 1995. Xylanases: from biology to biotechnology. *Biotechnol Gene Eng Rev.*, 13(12):101–131.
- [6] Zhang H, Li J, Wang J, et al., 2014, Determinants for the improved thermostability of a mesophilic family 11 xylanase predicted by computational methods. *Biotechnology for Biofuels.*, 7(1): 3–13.